

Unified parametrization of quark and lepton mixing matrices in tri-bimaximal pattern

Shi-Wen Li

School of Physics, Peking University, Beijing 100871, China

Bo-Qiang Ma*

School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China

Parametrization of the quark and lepton mixing matrices is the first attempt to understand the mixing of fermions. In this work, we parameterize the quark and lepton matrices with the help of quark-lepton complementarity (QLC) in a tri-bimaximal pattern of lepton mixing matrix. In this way, we combine the parametrization of the two matrices with each other. We apply this new parametrization to several physical quantities, and show its simplicity in the expression of, e.g., the Jarlskog parameter of CP violation.

PACS numbers: 14.60.Pq, 12.15.Ff, 14.60.Lm

I. INTRODUCTION

Mixing of different generations of fermions is one of the most important unsolved problems in particle physics, and has already troubled physicists for half a century. Quark mixing of two generations was first introduced by Cabibbo [1] decades ago as a unified explanation to the difference of the coupling constants in β decay and Λ decay in weak interaction. This idea was extended to the case of three generations by Kobayashi and Maskawa [2]. Parallely, neutrino oscillation as the lepton mixing was originally motivated by Pontecorvo [3] fifty years ago to understand the solar neutrino deficit, and further generalized by Maki, Nakagawa and Sakata [4]. Visually, these mixings could be regarded as a rotation from fermion mass eigenstates to flavor eigenstates. Therefore, this rotation could be expressed as a mixing matrix, namely the Cabibbo-Kobayashi-Maskawa (CKM) matrix V for quark sector and the Pontecorvo-Maki-Nakawaga-Sakata (PMNS) matrix U for lepton sector,

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}, \quad U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix}.$$

Both the CKM and PMNS matrices are unitary matrices, comprising three mixing angles θ_{ij} ($i, j = 1, 2, 3$) and one Dirac CP-violating phase δ , and could thus be rewritten as

$$\begin{pmatrix} c_{12}c_{31} & s_{12}c_{31} & s_{31}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{31}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{31}e^{i\delta} & s_{23}c_{31} \\ s_{12}s_{23} - c_{12}c_{23}s_{31}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{31}e^{i\delta} & c_{23}c_{31} \end{pmatrix} V = \begin{pmatrix} 1 - \frac{1}{2}\lambda^2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{1}{2}\lambda^2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \mathcal{O}(\lambda^4), \quad (2)$$

where $s_{ij} = \sin \theta_{ij}$ and $c_{ij} = \cos \theta_{ij}$. This expression was first introduced by Chau and Keung [5] in 1984, and was advocated by the Particle Data Group in 1996. If neutrinos are of Majorana type, there should be one additional diagonal matrix with two Majorana phases

$\text{diag}(e^{i\alpha_1/2}, e^{i\alpha_2/2}, 1)$ multiplied to the matrix above. However, the two Majorana phases do not affect neutrino oscillations, and we ignore the diagonal matrix here.

The similarity of the CKM and PMNS matrices arouses a natural question: what are their relations, e.g., the relations between their elements, or equivalently, mixing angles? Is there a unified comprehension of the two mixing matrices, just like the Cabibbo's way out to interpret different decays with the same coupling constant? Before the underlying grand unified theory of particle physics is discovered, phenomenological analyses play the main role in the correct direction. Amongst phenomenological analyses, to parameterize the CKM and PMNS matrices may be the first attempt. Parametrization provides us a numerical understanding of the hierarchy of different elements in the mixing matrix, and this is the purpose of our work.

For the CKM matrix, various particle physics experiments have precisely fixed its value, with the allowed ranges of the magnitudes of its elements reading [6]

$$\begin{pmatrix} 0.97383^{+0.00024}_{-0.00023} & 0.2272^{+0.0010}_{-0.0010} & 0.00396^{+0.00009}_{-0.00009} \\ 0.2271^{+0.0010}_{-0.0010} & 0.97296^{+0.00024}_{-0.00024} & 0.04221^{+0.00010}_{-0.00010} \\ 0.00814^{+0.00032}_{-0.00064} & 0.04161^{+0.00012}_{-0.00078} & 0.999100^{+0.000034}_{-0.000004} \end{pmatrix}. \quad (1)$$

We find that the CKM matrix is approximately the unit matrix with small deviations in the non-diagonal elements, and it could be parameterized by the standard Wolfenstein parametrization [7],

with $\lambda = 0.2272^{+0.0010}_{-0.0010}$ and $A = 0.818^{+0.007}_{-0.017}$ [6]. Nevertheless, the situation of the PMNS matrix is exactly opposite. This is because (1) the long-existed inaccuracy of neutrino experiments caused a delay in a practical parametrization of the PMNS matrix as clear as the CKM matrix; (2) almost all the elements in the PMNS matrix are of order $\mathcal{O}(1)$ except U_{e3} , which makes the direct application of the Wolfenstein-like perturbative expansion infeasible.

*Corresponding author. Email address: mabq@phy.pku.edu.cn

Whereas, circumstances have changed dramatically in recent years. Thanks to more and more precise measurements of neutrino oscillations, we have now a solid knowledge of the properties of neutrinos, aspects of their mass squared differences and mixing angles (although not as accurate as those of quarks). This helps us to remove the first obstacle mentioned above, and in this paper, we focus on the second one, i.e., how to parameterize the PMNS matrix with large elements? Moreover, we try to find its relation with the CKM matrix with the help of the so-called “quark-lepton complementarity” (i.e., the sum of corresponding mixing angles in quark and lepton sectors is about 45° , always referred as QLC in literature), and parameterize both of them.

The organization of this paper is: in Sec. II, we take the tri-bimaximal pattern as the basis of the PMNS matrix, but do not parameterize the PMNS matrix around it immediately. However, we use the QLC to obtain a new basis for the CKM matrix, and discuss its parametrization. In Sec. III, with the new parameterized CKM matrix, we utilize the QLC again, and the parametrization of the PMNS matrix around the tri-bimaximal pattern is consequently derived, with the same set of parameters in quark sector. Thus, we attain a unified parametrization of both the CKM and PMNS matrices. Several physical applications of the parametrization of the PMNS matrix are also studied in detail. After that, we talk about the relations between the two matrices and confirm the usefulness of the QLC when we consider the parametrization of the PMNS matrix as an independent one. Conclusions and discussions are summarized in Sec. IV.

II. PARAMETRIZATION OF THE CKM MATRIX

Parametrization of the CKM matrix has already been well established, i.e., the Wolfenstein parametrization in Eq. (2), which explicitly show the deviations of the non-diagonal elements from the unit matrix at different powers of the parameter λ . While, this method was presented long before the release of enormous experimental data about neutrinos, and its parameters were introduced, independently. Maybe a method of parametrization connecting the quark sector to the lepton sector is more effective at the epoch when fruitful neutrino experimental data are currently available. Therefore, in this section, we contemplate a new method for the parametrization of the CKM matrix, i.e., we parameterize it around a new basis, which is obtained from the tri-bimaximal pattern of the PMNS matrix by means of the QLC. Before continuing this procedure, let us outline the main results of lepton mixing (i.e., neutrino oscillations).

Nowadays, vast experimental data have confirmed the oscillations between different generations of neutrinos. For solar neutrino oscillation, a global analysis of the KamLAND experiments yielded $\Delta m_{21}^2 = 7.9^{+0.6}_{-0.5} \times 10^{-5} \text{ eV}^2$ and $\tan^2 \theta_{12} = 0.40^{+0.10}_{-0.07}$ [8]; as for SNO experiments, $\Delta m_{21}^2 = 8.0^{+0.6}_{-0.4} \times 10^{-5} \text{ eV}^2$ and $\theta_{12} =$

parameters	best fit	2σ	3σ
Δm_{21}^2	7.6	7.3 - 8.1	7.1 - 8.3
$ \Delta m_{31}^2 $	2.4	2.1 - 2.7	2.0 - 2.8
$\sin^2 \theta_{12}$	0.32	0.28 - 0.37	0.26 - 0.40
$\sin^2 \theta_{23}$	0.50	0.38 - 0.63	0.34 - 0.67
$\sin^2 \theta_{31}$	0.007	≤ 0.033	≤ 0.050

TABLE I: The three-flavor neutrino oscillation parameters from global data including solar, atmospheric, reactor (KamLAND and CHOOZ) and accelerator (K2K and MINOS) experiments. Δm_{21}^2 is in units of 10^{-5} eV^2 and $|\Delta m_{31}^2|$ of 10^{-3} eV^2 .

$(33.9^{+2.4}_{-2.2})^\circ$ [9]. Atmospheric neutrino oscillation experimental data by different groups are: $1.5 \times 10^{-3} \text{ eV}^2 < |\Delta m_{23}^2| < 3.4 \times 10^{-3} \text{ eV}^2$ and $\sin^2 2\theta_{23} > 0.92$ at 90% C.L. (Super-Kamiokande Collaboration [10]); $|\Delta m_{23}^2| = 2.74^{+0.44}_{-0.26} \times 10^{-3} \text{ eV}^2$ and $\sin^2 2\theta_{23} > 0.87$ at 68% C.L. (MINOS Collaboration [11]); $1.9 \times 10^{-3} \text{ eV}^2 < |\Delta m_{23}^2| < 3.5 \times 10^{-3} \text{ eV}^2$ at 90% C.L. when $\sin^2 2\theta_{23} = 1$, with the best fit value of $2.8 \times 10^{-3} \text{ eV}^2$ (K2K Collaboration [12]). On the other hand, the CHOOZ experiment [13] indicated that the upper bound of the mixing angle θ_{31} is pretty small. (See Table I [14] for a summary of the updated neutrino oscillation parameters and also Ref. [15] for an excellent illustration of the oscillation parameters.) According to the results of global analysis, the modulus of the PMNS matrix is summarized as (at 3σ C.L.) [16]

$$|U| = \begin{pmatrix} 0.77 - 0.86 & 0.50 - 0.63 & < 0.22 \\ 0.22 - 0.56 & 0.44 - 0.73 & 0.57 - 0.80 \\ 0.21 - 0.55 & 0.40 - 0.71 & 0.59 - 0.82 \end{pmatrix}, \quad (3)$$

Eq. (3) manifests the nontriviality of the non-diagonal elements in the PMNS matrix, and illuminates a distinct scheme for the parametrization of the PMNS matrix. We should not apply the method of the Wolfenstein parametrization of the CKM matrix directly to the PMNS matrix, so firstly we need to find another basis for the PMNS matrix. Otherwise, the parameter used for the expansion would be of order $\mathcal{O}(1)$, the expansion of the PMNS matrix then converges very slowly, and the hierarchy of different elements is also rather vague.

The matrix chosen as the basis for our intention must be close to the experimental constraints in Eq. (3). The most popular basis matrices are the “bimaximal” pattern (with $\sin \theta_{12} = \sin \theta_{23} = \sqrt{2}/2$) and the “tri-bimaximal” pattern (with $\sin \theta_{12} = \sqrt{3}/3$ and $\sin \theta_{23} = \sqrt{2}/2$). The parametrization of the PMNS matrix based on the bi-maximal pattern was discussed in [17]. In this paper, we concentrate on the parametrization of the PMNS matrix around the tri-bimaximal pattern.

Interestingly, the tri-bimaximal pattern was also intro-

duced by Wolfenstein [18] thirty years ago with the form

$$U = \begin{pmatrix} \sqrt{6}/3 & \sqrt{3}/3 & 0 \\ -\sqrt{6}/6 & \sqrt{3}/3 & \sqrt{2}/2 \\ \sqrt{6}/6 & -\sqrt{3}/3 & \sqrt{2}/2 \end{pmatrix}. \quad (4)$$

Recently, this special matrix has received dense discussions in literature [19]. Comparing Eq. (3) with Eq. (4), we find that the tri-bimaximal pattern agrees with the current experimental data fairly well, and the parametrization around it must be reasonable.

Parametrization of the PMNS matrix around the tri-bimaximal pattern was treated in several approaches [20]. Here, we do not repeat these approaches, but postpone this parametrization in the next section. Now, we want to see what happens to the CKM matrix if we connect the tri-bimaximal pattern with the QLC.

QLC is an appealing relation between the quark and lepton mixing angles,

$$\begin{aligned} \theta_{12}^{\text{CKM}} + \theta_{12}^{\text{PMNS}} &= 45^\circ, & \theta_{23}^{\text{CKM}} + \theta_{23}^{\text{PMNS}} &= 45^\circ, \\ \theta_{31}^{\text{CKM}} &\approx \theta_{31}^{\text{PMNS}} \approx 0^\circ. \end{aligned} \quad (5)$$

Below we use the superscripts CKM and PMNS to denote the parameters of the CKM and PMNS matrices, respectively. This relation was first introduced by Smirnov [21], and has been heavily studied in different sorts of models [22]. Eq. (5) is an extension for it.

Now that we have these empirical equations, the CKM and PMNS matrices are not independent of each other, and therefore we are enlightened to parameterize them in the same framework. With these numerical equations and the tri-bimaximal pattern of the PMNS matrix, we easily get the trigonometric functions of the mixing angles in the CKM matrix,

$$\begin{aligned} \sin \theta_{12}^{\text{CKM}} &= \frac{\sqrt{2}-1}{\sqrt{6}}, & \cos \theta_{12}^{\text{CKM}} &= \frac{\sqrt{2}+1}{\sqrt{6}}, \\ \sin \theta_{23}^{\text{CKM}} &= 0, & \cos \theta_{23}^{\text{CKM}} &= 1, \\ \sin \theta_{31}^{\text{CKM}} &= 0, & \cos \theta_{31}^{\text{CKM}} &= 1. \end{aligned}$$

The corresponding CKM matrix is thus fixed straightforwardly,

$$V = \begin{pmatrix} \frac{\sqrt{2}+1}{\sqrt{6}} & \frac{\sqrt{2}-1}{\sqrt{6}} & 0 \\ -\frac{\sqrt{2}-1}{\sqrt{6}} & \frac{\sqrt{2}+1}{\sqrt{6}} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (6)$$

Eq. (6), being the lowest order approximation of the CKM matrix, is not symmetric with $\theta_{12}^{\text{CKM}} = 9.7^\circ$, $\theta_{23}^{\text{CKM}} = 0^\circ$ and $\theta_{31}^{\text{CKM}} = 0^\circ$. Although this matrix is a little more complicated than the unit matrix, it is closer to reality. If we take the bimaximal pattern of the PMNS matrix with the QLC, we find that we get the unit matrix as the lowest order approximation of the CKM matrix. In the following, we just take Eq. (6) as the new basis for the CKM matrix, and parameterize the CKM matrix around it.

Comparing Eq. (6) with Eq. (1), we make an expansion of V in powers of λ (Attention, λ here is not the Wolfenstein one, i.e., $V_{us} = \lambda$),

$$V_{us} = \frac{\sqrt{2}-1}{\sqrt{6}} + \lambda,$$

where $\lambda = 0.0581_{-0.0010}^{+0.0010}$ measures the strength of deviation of V_{us} from the corresponding element in the matrix we just obtained in Eq. (6). Since $0 < \lambda < 0.1$, this expansion converges quickly. Similarly, we denote

$$V_{cb} = A\lambda,$$

and $A = 0.726_{-0.018}^{+0.013}$. As the CP violating phase is in V_{ub} , and $V_{cb} \sim 10|V_{ub}|$, we may set

$$V_{ub} = A\lambda^2(\rho - i\eta).$$

Thus, $\sqrt{\rho^2 + \eta^2} = 1.61_{-0.06}^{+0.05}$, and the values of ρ and η correspond to the Dirac CP-violating phase δ^{CKM} .

In summary, four parameters: λ , A , ρ and η are introduced here,

$$\begin{aligned} \lambda &= 0.0581_{-0.0010}^{+0.0010}, & A &= 0.726_{-0.018}^{+0.013}, \\ \sqrt{\rho^2 + \eta^2} &= 1.61_{-0.06}^{+0.05}, & 1.23 &\lesssim \eta/\rho \lesssim 4.70. \end{aligned}$$

where the ratio η/ρ is fixed by $\eta/\rho \simeq \tan \gamma \equiv -\tan \arg[(V_{ud}V_{ub}^*)/(V_{cd}V_{cb}^*)]$ ($\gamma = (63_{-12}^{+15})^\circ$) [6], and these four parameters characterize the CKM matrix completely.

Unitarity then determines the CKM matrix

$$\begin{aligned} V &= \begin{pmatrix} \frac{\sqrt{2}+1}{\sqrt{6}} & \frac{\sqrt{2}-1}{\sqrt{6}} & 0 \\ -\frac{\sqrt{2}-1}{\sqrt{6}} & \frac{\sqrt{2}+1}{\sqrt{6}} & 0 \\ 0 & 0 & 1 \end{pmatrix} + \lambda \begin{pmatrix} -(3-2\sqrt{2}) & 1 & 0 \\ -1 & -(3-2\sqrt{2}) & A \\ \frac{\sqrt{2}-1}{\sqrt{6}}A & -\frac{\sqrt{2}+1}{\sqrt{6}}A & 0 \end{pmatrix} \\ &+ \lambda^2 \begin{pmatrix} -(30\sqrt{3}-21\sqrt{6}) & 0 & (\rho-i\eta)A \\ \frac{\sqrt{2}-1}{2\sqrt{6}}A^2 & -(30\sqrt{3}-21\sqrt{6}) - \frac{\sqrt{2}+1}{2\sqrt{6}}A^2 & 0 \\ \left(1 - \frac{\sqrt{2}+1}{\sqrt{6}}(\rho+i\eta)\right)A & \left(3-2\sqrt{2} - \frac{\sqrt{2}-1}{\sqrt{6}}(\rho+i\eta)\right)A & -\frac{1}{2}A^2 \end{pmatrix} + \mathcal{O}(\lambda^3). \end{aligned} \quad (7)$$

We can extract the information of quark mixing from Eq. (7):

1. The expansion is reasonable in powers of λ and converges quickly.
2. The term of λ^0 is the new basis we have introduced in Eq. (6).
3. The term of λ^1 clearly shows the deviation of the CKM matrix from this new basis, and it contains only one parameter A except λ .
4. The term of λ^2 is the modification of higher order, and the effect of CP violation exists in this order. Since CP violation is stored in V_{ub} , the degree of CP violation in quark sector is of order λ^2 in our parametrization.

In this section, the CKM matrix is parameterized in a new way, i.e., around a new matrix, which is considered as the basis of the CKM matrix. The crucial input is the QLC. Following this idea, we make use of the QLC once more, and the parametrization of the PMNS matrix is then obtained.

III. PARAMETRIZATION OF THE PMNS MATRIX

Although many properties of neutrinos have been known from neutrino oscillation experiments, information is not enough to ensure the accurate ranges of the modulus of the elements in the PMNS matrix. Known from Eq. (3), the parametrization of the PMNS matrix is troublesome if we do not take a basis matrix into account. So we need a method to deal with it and make the parametrization feasible. Fortunately, the QLC connects the mixing angles of the two matrices which indicates that we may get the PMNS matrix from the CKM matrix. In this paper, we do not take the Wolfenstein parametrization of the CKM matrix as the perturbative expansion which leads to another parametrization of the PMNS matrix with the bimaximal pattern (see [23] for details), instead, we use Eq. (7) we have just obtained for

the CKM matrix. With the QLC, we find the trigonometric functions of the mixing angles in the PMNS matrix (to order of λ^2),

$$\begin{aligned}\sin \theta_{12}^{\text{PMNS}} &= \frac{\sqrt{3}}{3} - (2\sqrt{2} - 2)\lambda - (15\sqrt{6} - 21\sqrt{3})\lambda^2, \\ \cos \theta_{12}^{\text{PMNS}} &= \frac{\sqrt{6}}{3} + (2 - \sqrt{2})\lambda - (15\sqrt{6} - 21\sqrt{3})\lambda^2, \\ \sin \theta_{23}^{\text{PMNS}} &= \frac{\sqrt{2}}{2} \left(1 - A\lambda - \frac{1}{2}A^2\lambda^2 \right), \\ \cos \theta_{23}^{\text{PMNS}} &= \frac{\sqrt{2}}{2} \left(1 + A\lambda - \frac{1}{2}A^2\lambda^2 \right), \\ \sin \theta_{31}^{\text{PMNS}} &= A\lambda^2 \sqrt{\zeta^2 + \xi^2}, \\ \cos \theta_{31}^{\text{PMNS}} &= 1,\end{aligned}\tag{8}$$

where A and λ are the parameters we have defined for the CKM matrix, i.e., we describe the CKM and PMNS matrices with the same set of parameters. As we see in the parametrization of the CKM matrix, there are totally four parameters, but from Eq. (5) we only know two precise numerical equations. It implies that we have to introduce another two new parameters ζ and ξ to describe the PMNS matrix completely. From the fifth equation in Eq. (8), $\sin \theta_{31}^{\text{PMNS}} e^{-i\delta^{\text{PMNS}}} = A\lambda^2(\zeta - i\xi) = A\lambda^2 z^*$, where $z \equiv \zeta + i\xi$. According to the experimental data of neutrino oscillations, the value of $\sin \theta_{31}^{\text{PMNS}}$ has not been fixed yet, and what we have known is only its upper bound (see Table I). We find that $|U_{e3}| = \sin \theta_{31}^{\text{PMNS}} < 0.22$ at 3σ C.L. from Eq. (3). If 0.22 is taken for the value of $|U_{e3}|$, we find that $\sqrt{\zeta^2 + \xi^2} \sim 90$, which is unsuitable for the expansion of the PMNS matrix because the hierarchy of the first three orders would be indefinite. But if the value of $|U_{e3}|$ is smaller than 0.01, $\sqrt{\zeta^2 + \xi^2} \lesssim 4$ and $\sin \theta_{31}^{\text{PMNS}} e^{-i\delta^{\text{PMNS}}} = A\lambda^2 z^*$ is preferable. The value of $|U_{e3}|$ depends on the experimental data, hence we may also set $\sin \theta_{31}^{\text{PMNS}} e^{-i\delta^{\text{PMNS}}} = A\lambda(\zeta' - i\xi') = A\lambda z'^*$ ($\xi'/\zeta' = \xi/\zeta = \tan \delta^{\text{PMNS}}$), where $z'^* \equiv \zeta' + i\xi'$. $\sqrt{\zeta'^2 + \xi'^2} \sim 5$ if we take 0.22 for the value of $|U_{e3}|$, which means that $\sin \theta_{31}^{\text{PMNS}} e^{-i\delta^{\text{PMNS}}} = A\lambda z'^*$ would be better if the value of $|U_{e3}|$ is not small enough. Let us discuss the difference between the two cases.

Case 1: $\sin \theta_{31}^{\text{PMNS}} e^{-i\delta^{\text{PMNS}}} = A\lambda^2 z^*$.

With the trigonometric functions of the mixing angles in the PMNS matrix we have the parametrization of the PMNS matrix

$$\begin{aligned}U &= \begin{pmatrix} \frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} & 0 \\ -\frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} & \frac{\sqrt{2}}{2} \\ \frac{\sqrt{6}}{6} & -\frac{\sqrt{3}}{3} & \frac{\sqrt{2}}{2} \end{pmatrix} + \lambda \begin{pmatrix} 2 - \sqrt{2} & -(2\sqrt{2} - 2) & 0 \\ 2 - \sqrt{2} - \frac{1}{\sqrt{6}}A & \sqrt{2} - 1 + \frac{1}{\sqrt{3}}A & -\frac{\sqrt{2}}{2}A \\ -(2 - \sqrt{2}) - \frac{1}{\sqrt{6}}A & -(\sqrt{2} - 1) + \frac{1}{\sqrt{3}}A & \frac{\sqrt{2}}{2}A \end{pmatrix} \\ &+ \lambda^2 \begin{pmatrix} -(15\sqrt{6} - 21\sqrt{3}) & -(15\sqrt{6} - 21\sqrt{3}) & z^*A \\ 15\sqrt{3} - \frac{63}{\sqrt{6}} + \left(2 - \sqrt{2} - \frac{z}{\sqrt{3}}\right)A + \frac{1}{2\sqrt{6}}A^2 & -\left(15\sqrt{3} - \frac{63}{\sqrt{6}}\right) + \left(\sqrt{2} - 1 - \frac{z}{\sqrt{6}}\right)A - \frac{1}{2\sqrt{3}}A^2 & -\frac{\sqrt{2}}{4}A^2 \\ -\left(15\sqrt{3} - \frac{63}{\sqrt{6}}\right) + \left(2 - \sqrt{2} - \frac{z}{\sqrt{3}}\right)A - \frac{1}{2\sqrt{6}}A^2 & 15\sqrt{3} - \frac{63}{\sqrt{6}} + \left(\sqrt{2} - 1 - \frac{z}{\sqrt{6}}\right)A + \frac{1}{2\sqrt{3}}A^2 & -\frac{\sqrt{2}}{4}A^2 \end{pmatrix} \\ &+ \mathcal{O}(\lambda^3).\end{aligned}\tag{9}$$

Case 2: $\sin \theta_{31}^{\text{PMNS}} e^{-i\delta^{\text{PMNS}}} = A\lambda z'^*$.

In this case, the last two equations in Eq. (8) are replaced by (to order of λ^2)

$$\sin \theta_{31}^{\text{PMNS}} = A\lambda\sqrt{(\zeta'^2 + \xi'^2)}, \quad \cos \theta_{31}^{\text{PMNS}} = 1 - \frac{1}{2}A^2\lambda^2(\zeta'^2 + \xi'^2).$$

Repeating the former process, we have

$$U = \begin{pmatrix} \frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} & 0 \\ -\frac{\sqrt{6}}{6} & \frac{\sqrt{3}}{3} & \frac{\sqrt{2}}{2} \\ \frac{\sqrt{6}}{6} & -\frac{\sqrt{3}}{3} & \frac{\sqrt{2}}{2} \end{pmatrix} + \lambda \begin{pmatrix} 2 - \sqrt{2} & -(2\sqrt{2} - 2) & z'^*A \\ 2 - \sqrt{2} - \left(\frac{1}{\sqrt{6}} + \frac{z'}{\sqrt{3}}\right)A & \sqrt{2} - 1 + \left(\frac{1}{\sqrt{3}} - \frac{z'}{\sqrt{6}}\right)A & -\frac{\sqrt{2}}{2}A \\ -(2 - \sqrt{2}) - \left(\frac{1}{\sqrt{6}} + \frac{z'}{\sqrt{3}}\right)A & -(\sqrt{2} - 1) + \left(\frac{1}{\sqrt{3}} - \frac{z'}{\sqrt{6}}\right)A & \frac{\sqrt{2}}{2}A \end{pmatrix} \\ + \lambda^2 \begin{pmatrix} -(15\sqrt{6} - 21\sqrt{3}) - \frac{\sqrt{6}}{6}|z'|^2A^2 & -(15\sqrt{6} - 21\sqrt{3}) - \frac{\sqrt{3}}{6}|z'|^2A^2 & 0 \\ 15\sqrt{3} - \frac{63}{\sqrt{6}} + mA + pA^2 & -(15\sqrt{3} - \frac{63}{\sqrt{6}}) + nA - qA^2 & -\frac{\sqrt{2}}{4}(1 + |z'|^2)A^2 \\ -\left(15\sqrt{3} - \frac{63}{\sqrt{6}}\right) + mA - pA^2 & 15\sqrt{3} - \frac{63}{\sqrt{6}} + nA + qA^2 & -\frac{\sqrt{2}}{4}(1 + |z'|^2)A^2 \end{pmatrix} + \mathcal{O}(\lambda^3), \quad (10)$$

where

$$m = 2 - \sqrt{2} - \frac{z'}{\sqrt{2} + 1}, \quad n = \sqrt{2} - 1 + \frac{2z'}{2 + \sqrt{2}}, \quad p = \frac{1}{2\sqrt{6}} + \frac{z'}{\sqrt{3}}, \quad q = \frac{1}{2\sqrt{3}} - \frac{z'}{\sqrt{6}}.$$

Now we see the meanings of the expansions of U in Eq. (9) and Eq. (10):

1. The PMNS matrix U is expanded in powers of λ which makes the expansion converge quickly, the same as the CKM matrix V . So the parameter λ we choose for the CKM matrix is quite well.
2. The term of λ^0 is the zeroth order approximation, we find that it is just the tri-bimaximal pattern of the PMNS matrix, i.e., the matrix we want to take as the basis of the PMNS matrix. So the method of getting the expansion of the PMNS matrix from the expansion of the CKM matrix with the QLC is feasible.
3. The term of λ^1 shows the deviation of the PMNS matrix from the tri-bimaximal pattern. We find that the deviation is real and just relates with λ and A in Eq. (9). On the other hand, it is complex and relates with all four parameters in Eq. (10).
4. The term of λ^2 is the modification of higher order, and it is much more complicated in Eq. (10) than Eq. (9).
5. CP violation is described by U_{e3} in the PMNS matrix, and we find that the degree of CP violation in lepton sector is of order λ^2 in the parametrization of *case 1* but of order λ^1 in that of *case 2*. Since the choice of *case 1* or *case 2* depends on the value of $|U_{e3}|$, it implies that the size of CP violation is determined by the magnitude of $\sin \theta_{31}^{\text{PMNS}}$.

Now let us apply the parametrization of the PMNS matrix to some physical observables.

The amplitude for neutrinoless double beta decay ($0\nu\beta\beta$) is proportional to the quantity [6]

$$|\langle m_{\beta\beta} \rangle| \equiv \left| \sum_i U_{ei}^2 m_i \right|.$$

Since the two Majorana CP-violating phases are related here, we must take them into account.

In *case 1* (to order of λ^2),

$$|\langle m_{\beta\beta} \rangle_1| = \left| \frac{1}{3}(2m_1 e^{i\alpha_1} + m_2 e^{i\alpha_2}) + \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda(m_1 e^{i\alpha_1} - m_2 e^{i\alpha_2}) - (54 - 38\sqrt{2})\lambda^2(m_1 e^{i\alpha_1} - m_2 e^{i\alpha_2}) \right|.$$

We find that only m_1 and m_2 affect the value of $|\langle m_{\beta\beta} \rangle_1|$, m_3 vanishes here to order of λ^2 . Furthermore, there is only one parameter λ which in our parametrization shows up in $|\langle m_{\beta\beta} \rangle_1|$.

In *case 2* (to order of λ^2),

$$|\langle m_{\beta\beta} \rangle_2| = \left| \langle m_{\beta\beta} \rangle_1 + A^2\lambda^2 \left((\zeta' - i\xi')^2 m_3 - \frac{1}{3}(\zeta'^2 + \xi'^2)(2m_1 e^{i\alpha_1} + m_2 e^{i\alpha_2}) \right) \right|.$$

Apparently, $|\langle m_{\beta\beta} \rangle_2|$ is more complicated than $|\langle m_{\beta\beta} \rangle_1|$. Though the four parameters in our parametrization all appear in $|\langle m_{\beta\beta} \rangle_2|$, the effect of m_3 is still small, suppressed by λ^2 .

The rephasing-invariant Jarlskog parameter $\mathcal{J}$ [24] of CP violation is given by

$$\mathcal{J} \equiv \text{Im}(U_{e2}U_{\mu 3}U_{e3}^*U_{\mu 2}^*) = s_{12}s_{23}s_{31}c_{12}c_{23}c_{31}^2 \sin \delta.$$

An essential feature of $\mathcal{J}$ is that it is phase-convention independent which makes it very important when discussing CP violation. The parameter is expressed in a simple form in our parametrization (to order of λ^2). In *case 1*, we have

$$\mathcal{J} = \frac{\sqrt{2}}{6} A \lambda^2 \xi,$$

and in *case 2*, we have

$$\mathcal{J} = \left(\frac{\sqrt{2}}{6} - \frac{\sqrt{2}-1}{\sqrt{6}} \lambda \right) A \lambda \xi'.$$

If we rescale $\zeta' \rightarrow \lambda \zeta$ and $\xi' \rightarrow \lambda \xi$, we will find that $\mathcal{J}$ in *case 2* becomes equivalent to $\mathcal{J}$ in *case 1* (to order of λ^2). If we take $\sin^2 \theta_{31}^{\text{PMNS}} = 0.007$ as a preliminary estimate, $\sqrt{\zeta'^2 + \xi'^2} \sim 2$, then ζ' and ξ' are of order $\mathcal{O}(1)$, *case 2*

is preferable. However, if $\sin^2 \theta_{31}^{\text{PMNS}} = 0.0001$ or less, ζ and ξ are of order $\mathcal{O}(1)$, then *case 1* is preferred.

Leptonic unitarity triangles [25] may be constructed with the orthogonality of different pairs of rows or columns of the PMNS matrix. The matrix elements U_{e2} , U_{e3} and $U_{\mu 3}$ play important roles in the experiments and they have relatively simple forms in our parametrization. There are two unitarity triangles which contain the three matrix elements, the “ ν_2, ν_3 ” triangle and the “ ν_e, ν_μ ” triangle. Here we take the “ ν_2, ν_3 ” unitarity triangle as an example (for previous discussions see Bjorken and King in [20]).

The unitarity relation of the “ ν_2, ν_3 ” triangle is

$$U_{e2}U_{e3}^* + U_{\mu 2}U_{\mu 3}^* + U_{\tau 2}U_{\tau 3}^* = 0. \quad (11)$$

In *case 1* (to order of λ^2), we have

$$\begin{aligned} U_{e2}U_{e3}^* &= \frac{z}{\sqrt{3}} A \lambda^2, \\ U_{\mu 2}U_{\mu 3}^* &= \frac{1}{\sqrt{6}} + \left(1 - \frac{1}{\sqrt{2}}\right) \lambda - \left(\frac{45}{\sqrt{6}} - \frac{63}{\sqrt{12}} + \frac{z}{2\sqrt{3}} A + \frac{\sqrt{6}}{3} A^2\right) \lambda^2, \\ U_{\tau 2}U_{\tau 3}^* &= -\frac{1}{\sqrt{6}} - \left(1 - \frac{1}{\sqrt{2}}\right) \lambda + \left(\frac{45}{\sqrt{6}} - \frac{63}{\sqrt{12}} - \frac{z}{2\sqrt{3}} A + \frac{\sqrt{6}}{3} A^2\right) \lambda^2, \end{aligned}$$

and in *case 2* (to order of λ^2),

$$\begin{aligned} U_{e2}U_{e3}^* &= \frac{z'}{\sqrt{3}} A \lambda - \frac{2z'}{\sqrt{2}+1} A \lambda^2, \\ U_{\mu 2}U_{\mu 3}^* &= \frac{1}{\sqrt{6}} + \left(1 - \frac{1}{\sqrt{2}} - \frac{z'}{2\sqrt{3}} A\right) \lambda - \left(\frac{45}{\sqrt{6}} - \frac{63}{\sqrt{12}} - \frac{z'}{\sqrt{2}+1} A + \left(\frac{\sqrt{6}}{3} - \frac{z'}{\sqrt{3}} + \frac{|z'|^2}{2\sqrt{6}}\right) A^2\right) \lambda^2, \\ U_{\tau 2}U_{\tau 3}^* &= -\frac{1}{\sqrt{6}} - \left(1 - \frac{1}{\sqrt{2}} + \frac{z'}{2\sqrt{3}} A\right) \lambda + \left(\frac{45}{\sqrt{6}} - \frac{63}{\sqrt{12}} + \frac{z'}{\sqrt{2}+1} A + \left(\frac{\sqrt{6}}{3} - \frac{z'}{\sqrt{3}} + \frac{|z'|^2}{2\sqrt{6}}\right) A^2\right) \lambda^2. \end{aligned}$$

We find that the $U_{e2}U_{e3}^*$ side of the “ ν_2, ν_3 ” unitarity triangle is very small, suppressed by λ^2 in *case 1* and λ in *case 2*. This is because $|U_{e2}|$ is of $\mathcal{O}(1)$, and $|U_{e3}|$ is of $\mathcal{O}(\lambda^2)$ in *case 1* but $\mathcal{O}(\lambda)$ in *case 2*. Obviously, Eq. (11) has been satisfied in both cases (to order of λ^2). If neutrinos are of Majorana type, there should be one additional factor $e^{i\alpha_2/2}$ multiplied to every expression, but Eq. (11) still holds.

At last, we turn to neutrino oscillations. The probability of the oscillation from flavor ν_α to flavor ν_β , $P(\nu_\alpha \rightarrow \nu_\beta)$ is given by

$$\begin{aligned} P(\nu_\alpha \rightarrow \nu_\beta) &= \left| \sum_{i=1}^3 U_{\alpha i}^* e^{-im_i^2 \frac{L}{2E}} U_{\beta i} \right|^2 \\ &= \delta_{\alpha\beta} - 4 \sum_{i>j} \text{Re}(U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*) \sin^2 \Delta_{ij} + 2 \sum_{i>j} \text{Im}(U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*) \sin 2\Delta_{ij}, \end{aligned}$$

where $\alpha, \beta = e, \mu, \tau$, $\Delta_{ij} \equiv (m_i^2 - m_j^2)L/4E$, L is the oscillation length and E is the beam energy of neutrinos.

Let us take $P(\nu_e \rightarrow \nu_e)$ for example (for other oscillation probabilities see the Appendix). In *case 1* (to order of λ^2), we obtain

$$P(\nu_e \rightarrow \nu_e) = 1 - \left(\frac{8}{9} - \frac{16}{9}(\sqrt{6} - \sqrt{3})\lambda - 8(\sqrt{2} - 1)\lambda^2 \right) \sin^2 \Delta_{21},$$

and in *case 2* (to order of λ^2), we have

$$P(\nu_e \rightarrow \nu_e) = 1 - \left(\frac{8}{9} - \frac{16}{9}(\sqrt{6} - \sqrt{3})\lambda - \left(8(\sqrt{2} - 1) + \frac{16}{9}(\zeta'^2 + \xi'^2)A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\ - \frac{4}{3}(\zeta'^2 + \xi'^2)A^2 \lambda^2 (2 \sin^2 \Delta_{31} + \sin^2 \Delta_{32}).$$

Certainly we find that $\sum_{\beta} P(\nu_{\alpha} \rightarrow \nu_{\beta}) = 1$ to order of λ^2 , which means that the sum of the probability that a neutrino changes its flavor and the probability that it does not is unity.

Nevertheless, in despite of the CKM matrix and the QLC, we may consider Eq. (8) as an independent parametrization of the PMNS matrix based on the tri-bimaximal pattern. Then Eq. (9) and Eq. (10) are still obtained. According to the data in Table I, we find the ranges of the parameters in the PMNS matrix (at 3σ C.L.),

$$-0.069 < \lambda^{\text{PMNS}} < 0.079, \\ -0.17 < A^{\text{PMNS}} \lambda^{\text{PMNS}} < 0.16,$$

and $\sqrt{\zeta^2 + \xi^2} < 16$ in *case 1*, $\sqrt{\zeta'^2 + \xi'^2} < 1.3$ in *case 2*.

Without loss of generality, we set $A^{\text{PMNS}} > 0$, and we only acquire the upper bound of A^{PMNS} ,

$$A^{\text{PMNS}} < 2.5.$$

Clearly, the ranges of λ^{CKM} and A^{CKM} are included in the ranges of λ^{PMNS} and A^{PMNS} since the experiments for the PMNS matrix are not as accurate as those for the CKM matrix. It implies that the corresponding parameters of the two matrices may be correlated with each other to some extent when the parametrization of the PMNS matrix is independent. Let us see if there are any relations between the parameters of the CKM matrix and those of the PMNS matrix.

$$\sin(\theta_{12}^{\text{CKM}} + \theta_{12}^{\text{PMNS}}) = \frac{\sqrt{2}}{2} + (\sqrt{6} - \sqrt{3})(\lambda^{\text{CKM}} - \lambda^{\text{PMNS}}) - (30 - 21\sqrt{2})(\lambda^{\text{CKM}})^2 - (30\sqrt{2} - 42)(\lambda^{\text{PMNS}})^2 \\ + (9\sqrt{2} - 12)\lambda^{\text{CKM}}\lambda^{\text{PMNS}}, \\ \sin(\theta_{23}^{\text{CKM}} + \theta_{23}^{\text{PMNS}}) = \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}(A^{\text{CKM}}\lambda^{\text{CKM}} - A^{\text{PMNS}}\lambda^{\text{PMNS}}) - \frac{\sqrt{2}}{4}(A^{\text{CKM}}\lambda^{\text{CKM}} - A^{\text{PMNS}}\lambda^{\text{PMNS}})^2.$$

It is clear that the QLC is reobtained to zeroth order, and the two equations above coincide with the first two equations in Eq. (5) to order of λ^2 if $\lambda^{\text{CKM}} = \lambda^{\text{PMNS}}$ and $A^{\text{CKM}} = A^{\text{PMNS}}$. If the relation between θ_{31}^{CKM} and $\theta_{31}^{\text{PMNS}}$ is explicit, there should be an equation which links $\sqrt{\rho^2 + \eta^2}$ to $\sqrt{\zeta^2 + \xi^2}$, where the values of ρ and η correspond to δ^{CKM} , and the values of ζ and ξ to δ^{PMNS} . Thus, we state that the parameters in the PMNS matrix are not independent although the parametrization of the PMNS matrix is independent. The relations between the parameters of the CKM and PMNS matrices could be realized by the QLC. On another point of view, we may consider the QLC as the zeroth order approximation to the sum of the corresponding mixing angles of the CKM and PMNS matrices.

The expansion of the PMNS matrix has been derived directly based on the tri-bimaximal pattern of the PMNS matrix in another work (see the first paper in [23]). In that case, the parametrization is independent, and the expansion is reasonable in powers of λ ($|\lambda| < 0.1$ at 3σ C.L.). The degree of CP violation is of order λ^1 , the

Jarlskog parameter is combined of two terms, and the effect of m_3 is suppressed by λ^2 in the amplitude of neutrinoless double beta decay. It means that the expansion in that case is closer to *case 2* in this paper where the parameters in the CKM and PMNS matrices are not independent no matter the parametrization of the PMNS matrix is independent or not. The QLC plays an important role in the combination of the two matrices. Nevertheless, the magnitude of $\theta_{31}^{\text{PMNS}}$ is not explicit, hence the parametrization of the PMNS matrix is arbitrary to some extent and the choice between Eq. (9) and Eq. (10) depends on the experimental data of $|U_{e3}|$.

IV. SUMMARY

In this work, we introduce a new basis for the CKM matrix, which is closer to reality than the unit matrix. The unit matrix is symmetric with no mixing angles while the new basis matrix is not symmetric with one non-vanishing mixing angle. Taking the tri-bimaximal pat-

tern as the zeroth approximation for the PMNS matrix, the new basis of the CKM matrix is obtained via the QLC. We parameterize the CKM matrix around the new basis instead of the unit matrix, and find the expansion is suitable as a perturbative approximation.

The parametrization of the PMNS matrix is attained from the expansion of the CKM matrix with the QLC, then the two matrices are unified with the same set of parameters. Nevertheless, we get the expansion of the PMNS matrix, even if we consider parametrization in lepton sector as an independent one. In this case, however, the parameters of the CKM matrix are still associated with those of the PMNS matrix, since the QLC is reobtained when we combine the parameters of the two matrices.

Anyway, the expansions of the CKM and PMNS matrices are reasonable and converge quickly since the parameter λ is very small. The effect of CP violation is also seen from the parametrization of the two matrices.

Furthermore, some observable quantities are expressed in relatively simple forms or only related with one or two parameters in our parametrization. Finally, the two matrices are correlated with each other and useful for analyzing experimental data and phenomenology in particle physics.

Acknowledgments

We are grateful to Nan Li for valuable contributions and to Zhiqiang Guo for discussions. This work is partially supported by National Natural Science Foundation of China (Nos. 10721063, 10575003, 10528510), by the Key Grant Project of Chinese Ministry of Education (No. 305001), and by the Research Fund for the Doctoral Program of Higher Education (China).

APPENDIX

In the appendix, we list all other probabilities for neutrino oscillations (to order of λ^2).

In *case 1*,

$$\begin{aligned}
P(\nu_e \rightarrow \nu_\mu) &= \left(\frac{4}{9} - \frac{8}{9}(\sqrt{6} - \sqrt{3} - A)\lambda - \left(4(\sqrt{2} - 1) + \frac{16}{9} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{8}\zeta \right) A \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad + \frac{2\sqrt{2}}{3} \zeta A \lambda^2 (\sin^2 \Delta_{31} - \sin^2 \Delta_{32}) + \frac{\sqrt{2}}{3} \xi A \lambda^2 (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_e \rightarrow \nu_\tau) &= \left(\frac{4}{9} - \frac{8}{9}(\sqrt{6} - \sqrt{3} + A)\lambda - \left(4(\sqrt{2} - 1) - \frac{16}{9} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{8}\zeta \right) A \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad - \frac{2\sqrt{2}}{3} \zeta A \lambda^2 (\sin^2 \Delta_{31} - \sin^2 \Delta_{32}) - \frac{\sqrt{2}}{3} \xi A \lambda^2 (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_\mu \rightarrow \nu_e) &= \left(\frac{4}{9} - \frac{8}{9}(\sqrt{6} - \sqrt{3} - A)\lambda - \left(4(\sqrt{2} - 1) + \frac{16}{9} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{8}\zeta \right) A \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad + \frac{2\sqrt{2}}{3} \zeta A \lambda^2 (\sin^2 \Delta_{31} - \sin^2 \Delta_{32}) - \frac{\sqrt{2}}{3} \xi A \lambda^2 (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_\mu \rightarrow \nu_\mu) &= 1 - \left(\frac{2}{9} - \frac{4}{9}(\sqrt{6} - \sqrt{3} - 2A)\lambda - \left(2(\sqrt{2} - 1) + \frac{16}{9} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{8}\zeta \right) A - \frac{8}{9}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad - \left(\frac{1}{3} - \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda + \left(54 - 38\sqrt{2} + \frac{2\sqrt{2}}{3}\zeta A - \frac{4}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad - \left(\frac{2}{3} + \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda - \left(54 - 38\sqrt{2} + \frac{2\sqrt{2}}{3}\zeta A + \frac{8}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32}, \\
P(\nu_\mu \rightarrow \nu_\tau) &= - \left(\frac{2}{9} - \frac{4}{9}(\sqrt{6} - \sqrt{3})\lambda - \left(2\sqrt{2} - 2 + \frac{8}{9}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad + \left(\frac{1}{3} - \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda + \left(54 - 38\sqrt{2} - \frac{4}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad + \left(\frac{2}{3} + \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda - \left(54 - 38\sqrt{2} + \frac{8}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
&\quad + \frac{\sqrt{2}}{3} \xi A \lambda^2 (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}),
\end{aligned}$$

$$\begin{aligned}
P(\nu_\tau \rightarrow \nu_e) &= \left(\frac{4}{9} - \frac{8}{9}(\sqrt{6} - \sqrt{3} + A)\lambda - \left(4(\sqrt{2} - 1) - \frac{16}{9} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{8}\zeta \right) A \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad - \frac{2\sqrt{2}}{3} \zeta A \lambda^2 (\sin^2 \Delta_{31} - \sin^2 \Delta_{32}) + \frac{\sqrt{2}}{3} \xi A \lambda^2 (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_\tau \rightarrow \nu_\mu) &= - \left(\frac{2}{9} - \frac{4}{9}(\sqrt{6} - \sqrt{3})\lambda - \left(2\sqrt{2} - 2 + \frac{8}{9}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad + \left(\frac{1}{3} - \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda + \left(54 - 38\sqrt{2} - \frac{4}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad + \left(\frac{2}{3} + \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda - \left(54 - 38\sqrt{2} + \frac{8}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
&\quad - \frac{\sqrt{2}}{3} \xi A \lambda^2 (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_\tau \rightarrow \nu_\tau) &= 1 - \left(\frac{2}{9} - \frac{4}{9}(\sqrt{6} - \sqrt{3} + 2A)\lambda - \left(2(\sqrt{2} - 1) - \frac{16}{9} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{8}\zeta \right) A - \frac{8}{9}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad - \left(\frac{1}{3} - \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda + \left(54 - 38\sqrt{2} - \frac{2\sqrt{2}}{3}\zeta A - \frac{4}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad - \left(\frac{2}{3} + \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda - \left(54 - 38\sqrt{2} - \frac{2\sqrt{2}}{3}\zeta A + \frac{8}{3}A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32}.
\end{aligned}$$

In case 2,

$$\begin{aligned}
P(\nu_e \rightarrow \nu_\mu) &= \left(\frac{4}{9} - \frac{8}{9} \left(\sqrt{6} - \sqrt{3} - \left(1 + \frac{\sqrt{2}}{4}\zeta' \right) A \right) \lambda \right. \\
&\quad \left. - \left(4(\sqrt{2} - 1) + \frac{16}{9}(\sqrt{6} - \sqrt{3}) \left(1 - \frac{7\sqrt{2}}{8}\zeta' \right) A + \frac{8}{9}(\zeta'^2 + \xi'^2)A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad + \left(\frac{2\sqrt{2}}{3}\zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3})\zeta' A - \frac{4}{3}(\zeta'^2 + \xi'^2)A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad - \left(\frac{2\sqrt{2}}{3}\zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3})\zeta' A + \frac{2}{3}(\zeta'^2 + \xi'^2)A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
&\quad + \frac{\sqrt{2}}{3} \left(\xi' A \lambda - (\sqrt{6} - \sqrt{3})\xi' A \lambda^2 \right) (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_e \rightarrow \nu_\tau) &= \left(\frac{4}{9} - \frac{8}{9} \left(\sqrt{6} - \sqrt{3} + \left(1 + \frac{\sqrt{2}}{4}\zeta' \right) A \right) \lambda \right. \\
&\quad \left. - \left(4(\sqrt{2} - 1) - \frac{16}{9}(\sqrt{6} - \sqrt{3}) \left(1 - \frac{7\sqrt{2}}{8}\zeta' \right) A + \frac{8}{9}(\zeta'^2 + \xi'^2)A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad - \left(\frac{2\sqrt{2}}{3}\zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3})\zeta' A + \frac{4}{3}(\zeta'^2 + \xi'^2)A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad + \left(\frac{2\sqrt{2}}{3}\zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3})\zeta' A - \frac{2}{3}(\zeta'^2 + \xi'^2)A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
&\quad - \frac{\sqrt{2}}{3} \left(\xi' A \lambda - (\sqrt{6} - \sqrt{3})\xi' A \lambda^2 \right) (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}),
\end{aligned}$$

$$\begin{aligned}
P(\nu_\mu \rightarrow \nu_e) &= \left(\frac{4}{9} - \frac{8}{9} \left(\sqrt{6} - \sqrt{3} - \left(1 + \frac{\sqrt{2}}{4} \zeta' \right) A \right) \lambda \right. \\
&\quad - \left(4(\sqrt{2} - 1) + \frac{16}{9}(\sqrt{6} - \sqrt{3}) \left(1 - \frac{7\sqrt{2}}{8} \zeta' \right) A + \frac{8}{9}(\zeta'^2 + \xi'^2) A^2 \right) \lambda^2 \Big) \sin^2 \Delta_{21} \\
&\quad + \left(\frac{2\sqrt{2}}{3} \zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3}) \zeta' A - \frac{4}{3}(\zeta'^2 + \xi'^2) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad - \left(\frac{2\sqrt{2}}{3} \zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3}) \zeta' A + \frac{2}{3}(\zeta'^2 + \xi'^2) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
&\quad - \frac{\sqrt{2}}{3} \left(\xi' A \lambda - (\sqrt{6} - \sqrt{3}) \xi' A \lambda^2 \right) (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_\mu \rightarrow \nu_\mu) &= 1 - \left(\frac{2}{9} - \frac{4}{9} \left(\sqrt{6} - \sqrt{3} - \left(2 + \frac{\sqrt{2}}{2} \zeta' \right) A \right) \lambda \right. \\
&\quad - \left(2(\sqrt{2} - 1) + \frac{16}{9}(\sqrt{6} - \sqrt{3}) \left(1 - \frac{7\sqrt{2}}{8} \zeta' \right) A - \frac{8}{9} \left(1 + \frac{\sqrt{2}}{2} \zeta' - \frac{3}{8} \zeta'^2 + \frac{5}{8} \xi'^2 \right) A^2 \right) \lambda^2 \Big) \sin^2 \Delta_{21} \\
&\quad - \left(\frac{1}{3} - \frac{4}{3} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{2} \zeta' A \right) \lambda \right. \\
&\quad + \left(54 - 38\sqrt{2} - \frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3}) \zeta' A - \frac{4}{3} \left(1 + \sqrt{2} \zeta' - \frac{1}{4} \zeta'^2 - \frac{1}{4} \xi'^2 \right) A^2 \right) \lambda^2 \Big) \sin^2 \Delta_{31} \\
&\quad - \left(\frac{2}{3} + \frac{4}{3} \left(\sqrt{6} - \sqrt{3} - \frac{\sqrt{2}}{2} \zeta' A \right) \lambda \right. \\
&\quad - \left(54 - 38\sqrt{2} - \frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3}) \zeta' A + \frac{8}{3} \left(1 - \frac{\sqrt{2}}{2} \zeta' + \frac{1}{8} \zeta'^2 + \frac{1}{8} \xi'^2 \right) A^2 \right) \lambda^2 \Big) \sin^2 \Delta_{32}, \\
P(\nu_\mu \rightarrow \nu_\tau) &= - \left(\frac{2}{9} - \frac{4}{9}(\sqrt{6} - \sqrt{3}) \lambda - \left(2(\sqrt{2} - 1) + \frac{8}{9} \left(1 + \frac{\sqrt{2}}{2} \zeta' + \frac{5}{8} \zeta'^2 + \frac{13}{8} \xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
&\quad + \left(\frac{1}{3} - \frac{4}{3}(\sqrt{6} - \sqrt{3}) \lambda + \left(54 - 38\sqrt{2} - \frac{4}{3} \left(1 + \sqrt{2} \zeta' + \frac{3}{4} \zeta'^2 + \frac{3}{4} \xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad + \left(\frac{2}{3} + \frac{4}{3}(\sqrt{6} - \sqrt{3}) \lambda - \left(54 - 38\sqrt{2} + \frac{8}{3} \left(1 - \frac{\sqrt{2}}{2} \zeta' + \frac{3}{8} \zeta'^2 + \frac{3}{8} \xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
&\quad + \frac{\sqrt{2}}{3} \left(\xi' A \lambda - (\sqrt{6} - \sqrt{3}) \xi' A \lambda^2 \right) (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_\tau \rightarrow \nu_e) &= \left(\frac{4}{9} - \frac{8}{9} \left(\sqrt{6} - \sqrt{3} + \left(1 + \frac{\sqrt{2}}{4} \zeta' \right) A \right) \lambda \right. \\
&\quad - \left(4(\sqrt{2} - 1) - \frac{16}{9}(\sqrt{6} - \sqrt{3}) \left(1 - \frac{7\sqrt{2}}{8} \zeta' \right) A + \frac{8}{9}(\zeta'^2 + \xi'^2) A^2 \right) \lambda^2 \Big) \sin^2 \Delta_{21} \\
&\quad - \left(\frac{2\sqrt{2}}{3} \zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3}) \zeta' A + \frac{4}{3}(\zeta'^2 + \xi'^2) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
&\quad + \left(\frac{2\sqrt{2}}{3} \zeta' A \lambda - \left(\frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3}) \zeta' A - \frac{2}{3}(\zeta'^2 + \xi'^2) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
&\quad + \frac{\sqrt{2}}{3} \left(\xi' A \lambda - (\sqrt{6} - \sqrt{3}) \xi' A \lambda^2 \right) (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}),
\end{aligned}$$

$$\begin{aligned}
P(\nu_\tau \rightarrow \nu_\mu) = & - \left(\frac{2}{9} - \frac{4}{9}(\sqrt{6} - \sqrt{3})\lambda - \left(2(\sqrt{2} - 1) + \frac{8}{9} \left(1 + \frac{\sqrt{2}}{2}\zeta' + \frac{5}{8}\zeta'^2 + \frac{13}{8}\xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
& + \left(\frac{1}{3} - \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda + \left(54 - 38\sqrt{2} - \frac{4}{3} \left(1 + \sqrt{2}\zeta' + \frac{3}{4}\zeta'^2 + \frac{3}{4}\xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
& + \left(\frac{2}{3} + \frac{4}{3}(\sqrt{6} - \sqrt{3})\lambda - \left(54 - 38\sqrt{2} + \frac{8}{3} \left(1 - \frac{\sqrt{2}}{2}\zeta' + \frac{3}{8}\zeta'^2 + \frac{3}{8}\xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32} \\
& - \frac{\sqrt{2}}{3} \left(\xi' A \lambda - (\sqrt{6} - \sqrt{3})\xi' A \lambda^2 \right) (\sin 2\Delta_{21} - \sin 2\Delta_{31} + \sin 2\Delta_{32}), \\
P(\nu_\tau \rightarrow \nu_\tau) = & 1 - \left(\frac{2}{9} - \frac{4}{9} \left(\sqrt{6} - \sqrt{3} + \left(2 + \frac{\sqrt{2}}{2}\zeta' \right) A \right) \lambda \right. \\
& - \left(2(\sqrt{2} - 1) - \frac{16}{9}(\sqrt{6} - \sqrt{3}) \left(1 - \frac{7\sqrt{2}}{8}\zeta' \right) A - \frac{8}{9} \left(1 + \frac{\sqrt{2}}{2}\zeta' - \frac{3}{8}\zeta'^2 + \frac{5}{8}\xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{21} \\
& - \left(\frac{1}{3} - \frac{4}{3} \left(\sqrt{6} - \sqrt{3} + \frac{\sqrt{2}}{2}\zeta' A \right) \lambda \right. \\
& + \left(54 - 38\sqrt{2} + \frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3})\zeta' A - \frac{4}{3} \left(1 + \sqrt{2}\zeta' - \frac{1}{4}\zeta'^2 - \frac{1}{4}\xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{31} \\
& - \left(\frac{2}{3} + \frac{4}{3} \left(\sqrt{6} - \sqrt{3} + \frac{\sqrt{2}}{2}\zeta' A \right) \lambda \right. \\
& \left. - \left(54 - 38\sqrt{2} + \frac{2\sqrt{2}}{3}(\sqrt{6} - \sqrt{3})\zeta' A + \frac{8}{3} \left(1 - \frac{\sqrt{2}}{2}\zeta' + \frac{1}{8}\zeta'^2 + \frac{1}{8}\xi'^2 \right) A^2 \right) \lambda^2 \right) \sin^2 \Delta_{32}.
\end{aligned}$$

-
- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).
[2] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
[3] B. Pontecorvo, Sov. Phys. JETP **6**, 429 (1957).
[4] Z. Maki, M. Nakagawa, and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962).
[5] L.L. Chau and W.Y. Keung, Phys. Rev. Lett. **53**, 1802 (1984).
[6] W.-M. Yao *et al.*, J. Phys. **G33**, 1 (2006).
[7] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).
[8] KamLAND Collaboration, T. Araki *et al.*, Phys. Rev. Lett. **94**, 081801 (2005).
[9] SNO Collaboration, B. Aharmim *et al.*, Phys. Rev. **C72** 055502 (2005).
[10] Super-Kamiokande Collaboration, Y. Ashie *et al.*, Phys. Rev. **D71**, 112005 (2005).
[11] MINOS Collaboration, D.G. Michael *et al.*, Phys. Rev. Lett. **97**, 191801 (2006).
[12] K2K Collaboration, M.H. Ahn *et al.*, Phys. Rev. **D74**, 072003 (2006).
[13] CHOOZ Collaboration, M. Apollonio *et al.*, Phys. Lett. **B466**, 415 (1999); Eur. Phys. J. **C27** 331 (2003); Palo Verde Collaboration, F. Boehm *et al.*, Phys. Rev. **D64** 112001 (2001).
[14] M. Maltoni, T. Schwetz, M.A. Tortola and J.W.F. Valle, hep-ph/0405172v6, New J. Phys. **6**, 122 (2004).
[15] <http://hitoshi.berkeley.edu/neutrino/>.
[16] M.C. Gonzalez-Garcia, M. Maltoni, arXiv: 0704.1800 [hep-ph].
[17] W. Rodejohann, Phys. Rev. **D69**, 033005 (2004); N. Li and B.-Q. Ma, Phys. Lett. **B600**, 248 (2004).
[18] L. Wolfenstein, Phys. Rev. **D18**, 958 (1978).
[19] P.F. Harrison, D.H. Perkins and W.G. Scott, Phys. Lett. **B458**, 79 (1999); Phys. Lett. **B530**, 167 (2002); Z.Z. Xing, Phys. Lett. **B533**, 85 (2002); P.F. Harrison and W.G. Scott, Phys. Lett. **B535**, 163 (2002); Phys. Lett. **B557**, 76 (2003); X.G. He and A. Zee, Phys. Lett. **B560**, 87 (2003); C.I. Low and R.R. Volkas, Phys. Rev. **D68**, 033007 (2003); A. Zee, Phys. Rev. **D68**, 093002 (2003); G. Altarelli and F. Feruglio, Nucl. Phys. **B720**, 64 (2005); Nucl. Phys. **B741**, 215 (2006); F. Plentinger and W. Rodejohann, Phys. Lett. **B625**, 264 (2005); R. Friedberg and T.D. Lee, hep-ph/0606071; R.N. Mohapatra, S. Nasri and H.B. Yu, Phys. Lett. **B639**, 318 (2006); X.G. He, Y.Y. Keum and R.R. Volkas, JHEP **0604**, 039 (2006); G. Altarelli, F. Feruglio and Y. Lin, Nucl. Phys. **B775**, 31 (2007); F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, Nucl. Phys. **B775**, 120 (2007); Y. Koide, J. Phys. **G34**, 1653 (2007); I. de Medeiros Varzielas, S.F. King and G.G. Ross, Phys. Lett. **B644**, 153 (2007); Phys. Lett. **B648**, 201 (2007); C. Luhn, S. Nasri and P. Ramond, Phys. Lett. **B652** 27 (2007); C.S. Lam, Phys. Lett. **B656**, 193 (2007).
[20] N. Li and B.-Q. Ma, Phys. Rev. **D71**, 017302 (2005); J.D. Bjorken, P.F. Harrison and W.G. Scott, Phys. Rev. **D74**, 073012 (2006); S.F. King, Phys. Lett. **B659**, 244

- (2008); S. Pakvasa, W. Rodejohann and T.J. Weiler, Phys. Rev. Lett. **100**, 111801 (2008).
- [21] A.Y. Smirnov, hep-ph/0402264.
- [22] H. Minakata and A.Y. Smirnov, Phys. Rev. **D70**, 073009 (2004); M. Raidal, Phys. Rev. Lett. **93**, 161801 (2004); P.H. Frampton and R.N. Mohapatra, JHEP **0501**, 025 (2005); J. Ferrandis and S. Pakvasa, Phys. Rev. **D71**, 033004 (2005); S.K. Kang, C.S. Kim and J. Lee, Phys. Lett. **B619**, 129 (2005); S. Antusch, S.F. King and R.N. Mohapatra, Phys. Lett. **B618**, 150 (2005); M.A. Schmidt and A.Y. Smirnov, Phys. Rev. **D74**, 113003 (2006); F. Plentinger, G. Seidl and W. Winter, Phys. Rev. **D76**, 113003 (2007).
- [23] N. Li and B.-Q. Ma, Phys. Rev. **D71**, 097301 (2005).
- [24] C. Jarlskog, Phys. Rev. Lett. **55**, 1039 (1985); Z. Phys. **C29**, 491 (1985)
- [25] Y. Farzan and A.Y. Smirnov, Phys. Rev. **D65**, 113001 (2002); Z.Z. Xing and H. Zhang, Phys. Lett. **B618**, 131 (2005); H. Zhang and Z.Z. Xing, Eur. Phys. J. **C41**, 143 (2005).